

PAIN MANAGEMENT	
ADULT	PEDIATIRC (≤34KG)
BLS	
<ul style="list-style-type: none"> • Universal Protocol #601 • Medical (non-cardiac) <ul style="list-style-type: none"> ○ Position of comfort ○ Nothing by mouth • Cardiac chest pain – Chest Pain/Acute Coronary Syndrome Protocol #640 • Trauma – General Trauma Protocol #660 <ul style="list-style-type: none"> ○ Splint, ice, elevate as indicated 	<p>Same as Adult</p>
BLS Optional	
Pulse Oximetry – O ₂ administration per Airway Management Protocol #602	
ALS Standing Orders	
<ul style="list-style-type: none"> • For isolated orthopedic extremity injuries, dislocations, or burns without associated multi-system trauma and SBP > 90mmHg <ul style="list-style-type: none"> ○ Morphine up to 5 mg IV/IM ○ May repeat – not to exceed 10 mg total • For severe nausea associated with MS administration <ul style="list-style-type: none"> ○ Ondansetron 4 mg SLOW IV over 1 min or ○ 4 mg IM/PO ○ May repeat every 20 min not to exceed 12 mg total 	<ul style="list-style-type: none"> • For isolated extremity injuries, dislocations or burns without associated multi-system trauma and BP within normal limits for age <ul style="list-style-type: none"> ○ Morphine 0.1 mg/kg slow IV/IM ○ May repeat – not to exceed 5 mg total
Base Hospital Orders Only	
<ul style="list-style-type: none"> • Morphine administration with multisystem injuries or GCS < 15 • As needed 	<ul style="list-style-type: none"> • For severe nausea associated with MS administration <ul style="list-style-type: none"> ○ Ondansetron 4 mg SLOW IV over 1 min or ○ 4 mg IM/PO ○ Repeat per Base Hospital order • As needed
Notes	
<ul style="list-style-type: none"> • Document pain score prior to and post intervention(s) 	