

<b>ABDOMINAL/FLANK PAIN (MEDICAL)</b>	
<b>ADULT</b>	<b>PEDIATRIC (≤34 KG)</b>
<b>BLS</b>	
<ul style="list-style-type: none"> <li>• Universal Protocol #601</li> <li>• NPO</li> <li>• Monitor for orthostatic changes</li> <li>• If VS are unstable                             <ul style="list-style-type: none"> <li>○ Place supine</li> </ul> </li> </ul>	Same as Adult
<b>BLS Optional</b>	
Pulse Oximetry – Administer O <sub>2</sub> per Airway Management Protocol #602	
<b>ALS Standing Orders</b>	
<p>SBP &lt;100 mmHg or other signs of hypotension</p> <ul style="list-style-type: none"> <li>• <b>Normal Saline</b> IV up to 500 ml                             <ul style="list-style-type: none"> <li>○ May repeat once if hypotension persists</li> </ul> </li> <li>• Consider establishing secondary IV access</li> <li>• Consider 12-lead ECG</li> <li>• If shock is due to trauma refer to General Trauma Protocol #660</li> </ul>	<p>Signs of hypotension specific to age</p> <ul style="list-style-type: none"> <li>• <b>Normal Saline</b> IV 20 ml/kg                             <ul style="list-style-type: none"> <li>○ May repeat once if hypotension persists</li> </ul> </li> <li>• Consider establishing secondary IV access</li> <li>• If shock is due to trauma refer to General Trauma Protocol #660</li> </ul>
<b>Base Hospital Orders Only</b>	
<ul style="list-style-type: none"> <li>• <b>Morphine</b> for pain                             <ul style="list-style-type: none"> <li>○ Dosing per Pain Management Protocol #603</li> </ul> </li> <li>• As needed</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Morphine</b> for pain</li> <li>• <b>Ondansetron</b> for nausea                             <ul style="list-style-type: none"> <li>○ 4 mg slow IV push over 1 min</li> <li>○ 4 mg IM or PO</li> <li>○ Repeat per Base Hospital order</li> </ul> </li> <li>• As needed</li> </ul>
<b>Notes</b>	
<ul style="list-style-type: none"> <li>• Consider acute coronary syndrome with atypical presentation</li> </ul>	