

ALLERGIC REACTION/ANAPHYLAXIS	
ADULT	PEDIATRIC (≤34 KG)
BLS	
<ul style="list-style-type: none"> • Universal Protocol #601 • May assist with the administration of patient’s prescribed medication (i.e. Epi Auto injector, inhaler, etc.) 	<p>Same as Adult</p>
BLS Optional	
Pulse Oximetry – O ₂ administration per Airway Management Protocol #602	
ALS Standing Orders	
<p style="text-align: center;">Stable (Itching/rash)</p> <ul style="list-style-type: none"> • Contact Base for orders <p style="text-align: center;">Unstable (Dyspnea/Wheezing/Shock)</p> <ul style="list-style-type: none"> • Albuterol 2.5-5 mg via HHN/Mask/BVM with adjunct, over 5-10 min <ul style="list-style-type: none"> ○ repeat as needed • Epinephrine 1:1,000 0.01 mg/kg IM – not to exceed 0.5 mg <ul style="list-style-type: none"> ○ may repeat every 5 min, max 3 doses • Diphenhydramine 50 mg IV/IM <p style="text-align: center;">Extremis</p> <ul style="list-style-type: none"> • Epinephrine 1:1,000 0.01 mg/kg SL – not to exceed 0.5 mg <ul style="list-style-type: none"> ○ may repeat every 5 min, max 3 doses 	<p style="text-align: center;">Stable (Itching/rash)</p> <ul style="list-style-type: none"> • Contact Base for orders <p style="text-align: center;">Unstable (Dyspnea/Wheezing/Shock)</p> <ul style="list-style-type: none"> • Albuterol 2.5-5 mg via HHN/Mask/BVM with adjunct, over 5-10 min <ul style="list-style-type: none"> ○ repeat as needed • Epinephrine 1:1,000 0.01 mg/kg IM – not to exceed 0.3 mg <ul style="list-style-type: none"> ○ may repeat every 5 min, max 3 doses • Diphenhydramine 2 mg/kg IV/IM – not to exceed 50 mg <p style="text-align: center;">Extremis</p> <ul style="list-style-type: none"> • Epinephrine 1:1,000 0.01 mg/kg SL – not to exceed 0.3 mg <ul style="list-style-type: none"> ○ may repeat every 5 min, max 3 doses
Base Hospital Orders Only	
<p>Stable</p> <ul style="list-style-type: none"> • Diphenhydramine 50 mg IV/IM <p>Unresponsive to previous therapy</p> <ul style="list-style-type: none"> • Epinephrine 1:10,000 0.01 mg/kg slow IV titrated – not to exceed 0.5 mg <ul style="list-style-type: none"> • As needed 	<p>Stable</p> <ul style="list-style-type: none"> • Diphenhydramine 2 mg/kg IV/IM – not to exceed 50 mg <p>Unresponsive to previous therapy</p> <ul style="list-style-type: none"> • Epinephrine 1:10,000 0.01 mg/kg slow IV titrated – not to exceed 0.3 mg <ul style="list-style-type: none"> • As needed
Notes	