

ALLERGIC REACTION/ANAPHYLAXIS	
ADULT	PEDIATRIC (≤34 KG)
BLS	
<ul style="list-style-type: none"> • Universal Protocol #601 • Pulse Oximetry <ul style="list-style-type: none"> ○ O₂ administration per Airway Management Protocol #602 • May assist with the administration of patient’s prescribed medication (i.e. Epi Auto-injector, inhaler, etc.) 	Same as Adult
BLS Elective Skill	
<p style="text-align: center;">Unstable (Dyspnea/Wheezing/Shock)</p> <ul style="list-style-type: none"> • Adult 0.3 mg Epinephrine Auto-Injector administered in anterolateral thigh <ul style="list-style-type: none"> ○ May repeat, if indicated, every 5 min, max 3 doses 	<p style="text-align: center;">Unstable (Dyspnea/Wheezing/Shock)</p> <ul style="list-style-type: none"> • Pediatric (≥15 kg) 0.15 mg Epinephrine Auto-Injector administered in anterolateral thigh <ul style="list-style-type: none"> ○ May repeat, if indicated, every 5 min, max 3 doses
ALS Standing Orders	
<p style="text-align: center;">Stable (Itching/rash)</p> <ul style="list-style-type: none"> • Contact Base for orders <p style="text-align: center;">Unstable (Dyspnea/Wheezing/Shock)</p> <ul style="list-style-type: none"> • Albuterol 2.5-5 mg via HHN/Mask/BVM with adjunct, over 5-10 min <ul style="list-style-type: none"> ○ repeat as needed • Epinephrine 1:1,000 0.01 mg/kg IM – not to exceed 0.5 mg <ul style="list-style-type: none"> ○ may repeat every 5 min, max 3 doses • Diphenhydramine 50 mg IV/IM <p style="text-align: center;">Extremis</p> <ul style="list-style-type: none"> • Epinephrine 1:1,000 0.01 mg/kg SL – not to exceed 0.5 mg <ul style="list-style-type: none"> ○ may repeat every 5 min, max 3 doses 	<p style="text-align: center;">Stable (Itching/rash)</p> <ul style="list-style-type: none"> • Contact Base for orders <p style="text-align: center;">Unstable (Dyspnea/Wheezing/Shock)</p> <ul style="list-style-type: none"> • Albuterol 2.5-5 mg via HHN/Mask/BVM with adjunct, over 5-10 min <ul style="list-style-type: none"> ○ repeat as needed • Epinephrine 1:1,000 0.01 mg/kg IM – not to exceed 0.3 mg <ul style="list-style-type: none"> ○ may repeat every 5 min, max 3 doses • Diphenhydramine 2 mg/kg IV/IM – not to exceed 50 mg <p style="text-align: center;">Extremis</p> <ul style="list-style-type: none"> • Epinephrine 1:1,000 0.01 mg/kg SL – not to exceed 0.3 mg <ul style="list-style-type: none"> ○ may repeat every 5 min, max 3 doses
Base Hospital Orders Only	
<p style="text-align: center;">Stable</p> <ul style="list-style-type: none"> • Diphenhydramine 50 mg IV/IM <p>Unresponsive to previous therapy</p> <ul style="list-style-type: none"> • Epinephrine 1:10,000 0.01 mg/kg slow IV titrated – not to exceed 0.5 mg • As needed 	<p style="text-align: center;">Stable</p> <ul style="list-style-type: none"> • Diphenhydramine 2 mg/kg IV/IM – not to exceed 50 mg <p>Unresponsive to previous therapy</p> <ul style="list-style-type: none"> • Epinephrine 1:10,000 0.01 mg/kg slow IV titrated – not to exceed 0.3 mg • As needed

Notes
<ul style="list-style-type: none">• Auto-injector injection site should be exposed and cleansed with aseptic technique prior to injection• Follow manufacture's instructions when using Epinephrine auto-injector