

<b>SEIZURE (ACTIVE)</b>	
<b>ADULT</b>	<b>PEDIATRIC (≤34KG)</b>
<b>BLS</b>	
<ul style="list-style-type: none"> <li>• Universal Protocol #601</li> <li>• Prevent patient from injuring themselves – Do not restrain</li> </ul>	Same as Adults
<b>BLS Optional</b>	
Pulse Oximetry – O <sub>2</sub> administration per Airway Management Protocol #602	
<b>ALS Standing Orders</b>	
<ul style="list-style-type: none"> <li>• <b>Midazolam</b> <ul style="list-style-type: none"> <li>○ Up to 2 mg slow IV or</li> <li>○ 5 mg IM/IN (2.5 mg each nostril)</li> <li>○ May repeat once after 10 min</li> </ul> </li> <li>• Obtain blood sugar level</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Midazolam</b> <ul style="list-style-type: none"> <li>○ 0.1 mg/kg slow IV not to exceed 2mg</li> <li>○ 0.1 mg/kg IM/IN (max 0.3 mL each nostril)</li> </ul> </li> <li>• Obtain blood sugar level</li> </ul>
<b>Base Hospital Orders Only</b>	
<ul style="list-style-type: none"> <li>• As needed</li> </ul>	<ul style="list-style-type: none"> <li>• Repeat doses of <b>Midazolam</b></li> <li>• As needed</li> </ul>
<b>Notes</b>	