

<b>INTRANASAL MEDICATION</b>	
<b>ADULT</b>	<b>PEDIATRIC (≤34KG)</b>
<b>BLS</b>	
Universal Protocol #601	
<b>BLS Optional</b>	
Pulse Oximetry – O <sub>2</sub> administration per Airway Management Protocol #602	
<b>ALS Standing Orders</b>	
<ul style="list-style-type: none"> <li>• Place patient in supine/recumbent position</li> <li>• If in seated position it may be necessary to compress nares after administration</li> <li>• Draw up appropriate medication and appropriate amount into one or two syringes                             <ul style="list-style-type: none"> <li>○ Remove needle(s) and apply MAD</li> <li>○ Administer medication briskly to atomize medication</li> <li>○ Up to 1 mL maximum volume per nostril</li> <li>○ Evaluate and document response to medication</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>• Same as Adult</li> <li>• Up to 0.3 mL maximum volume per nostril</li> </ul>
<b>Base Hospital Orders Only</b>	
As needed	
<b>Notes</b>	
<ul style="list-style-type: none"> <li>• Contraindications                             <ul style="list-style-type: none"> <li>○ Epistaxis</li> <li>○ Nasal Trauma</li> </ul> </li> <li>• May not be effective in the case of:                             <ul style="list-style-type: none"> <li>○ Severe hypotension – preventing adequate absorption</li> <li>○ Recent use of nasal vasoconstriction medication</li> <li>○ Nasal septal abnormalities</li> <li>○ Nasal congestion or discharge</li> </ul> </li> </ul>	