

<b>SEIZURE (ACTIVE)</b>	
<b>ADULT</b>	<b>PEDIATRIC (≤34KG)</b>
<b>BLS</b>	
<ul style="list-style-type: none"> <li>• Universal Protocol #601</li> <li>• Pulse Oximetry                             <ul style="list-style-type: none"> <li>○ O<sub>2</sub> administration per Airway Management Protocol #602</li> </ul> </li> <li>• Prevent patient from injuring themselves – Do not restrain</li> </ul>	<p>Same as Adults</p>
<b>BLS Elective Skills</b>	
Obtain Blood Sugar Level – if <60 mg/dL see Altered Mental Status Protocol #612	
<b>ALS Standing Orders</b>	
<ul style="list-style-type: none"> <li>• <b>Midazolam</b> <ul style="list-style-type: none"> <li>○ Up to 2 mg slow IV or</li> <li>○ 5 mg IM/IN (split between nares)</li> <li>○ May repeat once after 10 min</li> </ul> </li> <li>• Obtain blood sugar level</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Midazolam</b> <ul style="list-style-type: none"> <li>○ 0.1 mg/kg slow IV not to exceed 2mg</li> <li>○ 0.1 mg/kg IM/IN (split between nares)</li> </ul> </li> <li>• Obtain blood sugar level</li> </ul>
<b>Base Hospital Orders Only</b>	
<ul style="list-style-type: none"> <li>• As needed</li> </ul>	<ul style="list-style-type: none"> <li>• Repeat doses of <b>Midazolam</b></li> <li>• As needed</li> </ul>
<b>Notes</b>	